

Published in final edited form as:

Curr Opin Neurobiol. 2011 October ; 21(5): 693–700. doi:10.1016/j.conb.2011.06.010.

New methods for localizing and manipulating neuronal dynamics in behaving animals

Michale S. Fee¹ and Michael A. Long²

¹McGovern Institute for Brain Research, Department of Brain and Cognitive Sciences, Massachusetts Institute of Technology, 77 Massachusetts Avenue, Cambridge, MA 02139 USA

²Departments of Otolaryngology and Physiology and Neuroscience, NYU School of Medicine, 522 First Ave., New York, NY 10016 USA

Abstract

Where are the ‘prime movers’ that control behavior? Which circuits in the brain control the order in which individual motor gestures of a learned behavior are generated, and the speed at which they progress? Here we describe two techniques recently applied to localizing and characterizing the circuitry underlying the generation of vocal sequences in the songbird. The first utilizes small, localized, temperature changes in the brain to perturb the speed of neural dynamics. The second utilizes intracellular manipulation of membrane potential in the freely behaving animal to perturb the dynamics within a single neuron. Both of these techniques are broadly applicable in behaving animals to test hypotheses about the biophysical and circuit dynamics that allow neural circuits to march from one state to the next.

Nearly everything humans learn to do requires the execution of a complex sequence of motor gestures produced in a precise temporal order and with precise timing. Much is now known about how neurons in motor areas spike in relation to individual motor gestures, or even sequences of gestures [1-6], for example, directional tuning of neurons in primate motor cortex [7-8] or sequence selectivity of neurons in supplementary motor cortex [9-10]. These are examples of how neuroscientists have begun to decipher how neurons in the brain fire in relation to — or *code for* — complex motor behaviors. But where are the ‘prime movers’ that drive complex learned behaviors — those circuits that generate the temporal structure of learned behaviors? What controls the order in which states of activity arise and the speed at which they progress? We need to go beyond a description of how neurons fire in relation to behavior to address the circuit and biophysical dynamics that underlie how one state of a neural system is transformed into the next state in time.

In some simple nervous systems, it has been possible to identify small circuits of neurons containing the biophysical dynamics that generate the temporal pattern of a behavior. For example, Stent and colleagues [11**] were able to classify subsets of neurons in the segmental ganglia of the leech as being part of an ‘oscillator network’ because manipulation of their membrane potential altered the timing of the rhythmic swimming behavior. In contrast, manipulating other ‘follower’ neurons had no effect on timing because their activity is simply driven by the oscillator network. Here we say that the dynamics that

© 2011 Elsevier Ltd. All rights reserved.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

underlie the swimming behavior are located entirely within the oscillator network, even though the follower neurons also exhibit time-dependent activity necessary for swimming.

Addressing these questions for more complex behaviors in large brains with many more neurons is more difficult. Lesions or pharmacological inactivation can tell us whether a brain region is necessary for the expression of a particular behavior. Electrophysiological recordings can tell us whether activity in a brain area is correlated with a behavior. Electrical stimulation can tell us whether activation of a region (or the axons passing into or through that region) can disrupt or elicit a behavior [1,12], but none of these approaches can reveal whether the biophysical dynamics within a region are actively involved in timing a behavior.

Here we describe two techniques that have recently been applied to the question of the origin of biophysical dynamics in neural circuits: Localized brain cooling and the intracellular manipulation of neuron membrane potential. Using these techniques, hypotheses can be tested about the dynamical origin of temporal structure in neural circuits that could not be tested using other techniques. We will focus on the application of these techniques to examine the origin of song timing in a songbird, an excellent model organism in which to address questions about the neuronal dynamics underlying complex learned sequences in the vertebrate brain. The song is mediated by a discrete set of motor nuclei, known as the song motor pathway, that are necessary for the production of adult song (Figure 1A) [13-17]. One nucleus, known as HVC (used as a proper name), is a cortical premotor brain region containing neurons that generate a single burst of spikes at a particular moment in the song. As a population, these neurons form a sparse sequential representation of time, or temporal order, in the song [18] (Figure 1B).

Using temperature to localize neuronal dynamics

Where are the dynamics that underlie the generation of sparse sequential bursts in HVC (Figure 2A)? One possibility is that the dynamics are contained entirely within HVC (Figure 2B). For example, one could envision a chain of activity in which each neuron is activated by a previous group of neurons and in turn activates the next group. An alternative possibility is that the bursting in HVC is directly driven by synaptic inputs from upstream brain areas, such as the thalamic nucleus Uva [16-17,19]. In the latter case, the timing of HVC activity could be controlled entirely by circuitry upstream of HVC (Figure 2C).

These two models give very different predictions for what would happen if we could slow down the biophysical processes in HVC. Imagine we could inject some sort of 'neuronal molasses' into HVC that doesn't inactivate the circuit, but just makes all neuronal processes move a little more slowly. If the circuitry and biophysical processes that control song timing reside entirely within HVC, then the time it takes each neuron to activate the next neuron in this sequence should increase. There should therefore be fewer bursts per unit time, and the song should slow down (Figure 2B).

The prediction is quite different if the burst sequence in HVC is driven by timing inputs from upstream of HVC. Slowing HVC could increase the duration of HVC bursts and would increase the time it takes HVC neurons to respond to this upstream input, but this increased latency would be the same for every HVC neuron. Thus, there would be the same number of bursts per unit time, and the song would not slow down (Figure 2C).

Temperature change is a useful manipulation to distinguish hypotheses like these about the role of specific circuits in timing of a behavior. Because it affects many neuronal processes such as synaptic transmission, axonal conduction velocity, and spiking properties [20-24], temperature change can be used very generally, even without detailed knowledge of the underlying circuit or neuronal properties. Although large decreases in temperature ($\Delta T >$

30°C) can result in inactivation [25-26], slight cooling ($\Delta T < 8^\circ\text{C}$) can slow circuit activity. Indeed, there are several examples now in which small temperature changes in simple central pattern generators have been shown to produce substantial changes in oscillatory cycle time [27, 28**,29] but otherwise leave the circuit and behavioral function intact [28**, 30*,31-32].

Temperature changes can be applied locally to distinguish the influence of even nearby brain areas, such as nucleus RA and HVC in the song motor pathway. The combination of thermal diffusion and local perfusion of the brain with a warm blood results in a localized effect of brain cooling. In birds, and probably mammals, the temperature change falls off from a cooling surface with a length constant of about 1.0-1.5mm [33]. Local cooling can be applied at the surface of the brain (for example, neocortex), or to structures deep within the brain using insulated probes. Using this approach, we developed a miniature device using small solid-state electronic heat pumps to provide local cooling of specific brain regions in the freely-behaving, singing zebra finches [33-34](Figure 3A,B). We found that bilateral cooling of HVC led to a nearly uniform slowing, or stretching, of the song by about 3% per degree of temperature change (Figure 3C). This stretch of the song occurred at all timescales: detailed acoustic structure within song syllables, the interval between syllable onsets, and the interval between song motifs. This finding provides strong evidence against models in which the timing of song vocalizations, on any timescale, is controlled by circuitry outside of HVC, and suggest that HVC may function as the 'high vocal clock' of song motor sequences.

Local temperature changes can be especially powerful when used to distinguish the roles of multiple interacting brain areas. If cooling one of these areas produces a slowing of the behavior while another does not, then strong conclusions can be drawn about the lack of involvement of the area in which cooling has no effect. For example, while cooling HVC causes a slowing of song structure, temperature changes in RA, the structure in the motor pathway to which HVC projects, did not result in a measurable change in song timing [33]. Taken together, these results suggest that biophysical dynamics within RA play little role in the control of song timing.

Observing an effect on timing from cooling a brain area may provide some evidence that timing originates in that area, however one must be careful to rule out other possible interpretations. For example, consider a situation in which the dynamics in one circuit (area B) are strongly sensitive to the amount of tonic 'activation' or perhaps neuromodulatory input from an upstream area (area A). Because cooling can substantially affect the firing rate of tonically active neurons (see ref [33] for an example), then cooling upstream area A could affect the temporal patterns in the downstream circuits, even though biophysical dynamics in area A are not directly involved in generating temporal patterns of activity. As with any technique, combining cooling with other approaches, such as electrophysiology, will be important to dissect neural circuit dynamics.

Manipulating intracellular membrane potential during behavior to test circuit models

Once HVC was localized as a principal locus of timing in the song pathway, it was then possible to address the detailed circuit mechanisms that underlie the sparse sequential activation of neurons in this area. Two major classes of models have been proposed to explain sequence generation within neural circuits. Groups of neurons may be linked together with specific local connections such that activity can propagate, wave-like, through a synaptically-connected 'chain' of neurons (Figure 4A, chain model) [35-39]. Alternatively, sequences could be generated in the absence of overt feed-forward connections between

neurons. For example, a population of neurons could exhibit a gradual and global ramping-up of membrane potential such that the most excitable neurons reach threshold first, while the least excitable neurons reach threshold later (Figure 4B, ramp-to-threshold model). Such global modulation of membrane potential is thought to underlie hippocampal theta sequences during theta activity [40] and perhaps replay sequences during sharp-wave activity [41-42]. In these models, the diversity of neuronal excitability (which controls the time at which each neuron spikes in the sequence) could be a function of intrinsic cellular excitability [43-44, 45*], or of recurrent network excitation [41].

Slow subthreshold inhibitory dynamics on the timescale of song syllables (~100 ms) has been observed in HVC *in vitro* [46], raising the possibility that such a ramp-to-threshold circuit could control burst timing in HVC. Note that any 'ramping' mechanism produced by biophysical dynamics within HVC would also likely be slowed by HVC cooling, so that the ramp-to-threshold model would predict a slowing of the song sequence by HVC cooling. Thus, the ramp-to-threshold model and the chain model described above would be very difficult to distinguish based on evidence from spike data or cooling experiments.

However, these two models yield very different predictions for the pattern of subthreshold activity that one would observe during singing. Specifically, in the ramp-to-threshold model, the membrane potential of each HVC neuron should exhibit a slow depolarizing ramp in the hundreds of milliseconds prior to burst onset, whereas in the chain model, each burst should be preceded by a fast depolarizing input from previously active neurons. Also, intracellular injection of hyperpolarizing current would be expected to make the recorded neuron reach threshold later, delaying the onset of the burst. Likewise, injection of depolarizing current would be expected to advance the time of the burst.

In order to test these predictions of these two circuit models, we adopted an approach recently introduced for intracellular recordings in the freely moving rat [47*,48] to carry out intracellular recordings in unrestrained, singing zebra finches. We developed a small (1.6 gram) microdrive and miniature headstage preamplifier for use with sharp glass microelectrodes (Figure 4C). With this device we were able to record synaptic inputs as well as spiking outputs from antidromically-identified premotor neurons in HVC (Figure 4D). The results were clearly consistent with the predictions of the chain model – in no cases did HVC neurons exhibit a ramping-up of membrane potential in the hundreds of milliseconds before burst onset. Indeed, bursts were preceded by a rapidly rising depolarizing potential within 5-10ms prior to burst onset, consistent with the idea that neurons are being driven by input from a previous 'link' in a chain of neurons. Furthermore, hyperpolarizing or depolarizing current injection did not change the time of the burst onset by more than a few milliseconds (Figure 4E), inconsistent with the idea that the time at which a neuron bursts is controlled by a slow ramping up to spike threshold.

Summary and Conclusions

Here we describe two approaches recently applied to localizing and characterizing the neuronal dynamics underlying the generation of complex behavioral sequences in behaving animals. While we have framed these ideas in terms of the circuits underlying behavior, the question of neuronal dynamics is pervasive in all aspects of brain function that involve time-dependent computations [49]. The tools we describe here should be broadly applicable to the study of brain dynamics underlying temporal selectivity [50*], changes in attention [51*], motor planning [52] and the generation of theta rhythm and phase precession important for hippocampal function [41,53**,54]. Other potential applications include examining the rich dynamics underlying the control of intracortical information flow [55*] and spontaneous

brain states [56*], or even localizing the dynamics underlying pathological states of brain activity such as Parkinsonian tremor [57*].

The tools we review here are complementary to recent approaches allowing optical control of activity in specific sets of neurons [58-66] or the targeting of ligand-gated channels to genetically specified sets of neurons in a circuit [67-68]. [69-70,71**]. These approaches will primarily be useful for studying neural circuits in which neuronal populations can be targeted with genetic tools, and about which enough is understood to formulate hypotheses about the function of specific circuit elements. In contrast, the broad effectiveness of localized brain cooling across neuronal subtypes and biophysical processes within a circuit makes this approach perfectly suited to the initial stage of cracking circuit dynamics – localizing, within a set of interconnected brain regions, where the neuronal dynamics are that underlie a behavior or a computation. Intracellular recording and current injection are similarly broadly applicable to address detailed questions about the dynamics of neuronal circuits in behaving animals.

References

1. Graziano MS, Aflalo TN. Mapping behavioral repertoire onto the cortex. *Neuron*. 2007; 56:239–251. [PubMed: 17964243]
2. Tanji J. Sequential organization of multiple movements: involvement of cortical motor areas. *Annu Rev Neurosci*. 2001; 24:631–651. [PubMed: 11520914]
3. Dombek DA, Graziano MS, Tank DW. Functional clustering of neurons in motor cortex determined by cellular resolution imaging in awake behaving mice. *J Neurosci*. 2009; 29:13751–13760. [PubMed: 19889987]
4. Churchland MM, Shenoy KV. Temporal complexity and heterogeneity of single-neuron activity in premotor and motor cortex. *J Neurophysiol*. 2007; 97:4235–4257. [PubMed: 17376854]
5. Hatsopoulos NG, Xu Q, Amit Y. Encoding of movement fragments in the motor cortex. *J Neurosci*. 2007; 27:5105–5114. [PubMed: 17494696]
6. Rizzolatti G, Luppino G. The cortical motor system. *Neuron*. 2001; 31:889–901. [PubMed: 11580891]
7. Georgopoulos AP, Schwartz AB, Kettner RE. Neuronal population coding of movement direction. *Science*. 1986; 233:1416–1419. [PubMed: 3749885]
8. Schwartz AB. Direct cortical representation of drawing. *Science*. 1994; 265:540–542. [PubMed: 8036499]
9. Shima K, Tanji J. Neuronal activity in the supplementary and presupplementary motor areas for temporal organization of multiple movements. *J Neurophysiol*. 2000; 84:2148–2160. [PubMed: 11024102]
10. Mushiake H, Inase M, Tanji J. Selective coding of motor sequence in the supplementary motor area of the monkey cerebral cortex. *Exp Brain Res*. 1990; 82:208–210. [PubMed: 2257906]
- **11. Stent GS, Kristan WB Jr, Friesen WO, Ort CA, Poon M, Calabrese RL. Neuronal generation of the leech swimming movement. *Science*. 1978; 200:1348–1357. This paper describes the ‘oscillator network’ in the segmental ganglia of the leech that underlies rhythmic swimming. The experiments involved intracellular current injection into single neurons in a fictive swimming preparation to demonstrate that manipulation of some alters the timing of the behavior, while manipulation of other neurons does not. By combining these findings with detailed functional circuit connectivity, the authors describe a fairly complete model of the underlying circuitry. [PubMed: 663615]
12. Vu ET, Mazurek ME, Kuo YC. Identification of a forebrain motor programming network for the learned song of zebra finches. *J Neurosci*. 1994; 14:6924–6934. [PubMed: 7965088]
13. Simpson HB, Vicario DS. Brain pathways for learned and unlearned vocalizations differ in zebra finches. *J Neurosci*. 1990; 10:1541–1556. [PubMed: 2332796]
14. Aronov D, Andalman AS, Fee MS. A specialized forebrain circuit for vocal babbling in the juvenile songbird. *Science*. 2008; 320:630–634. [PubMed: 18451295]

15. Nottebohm F, Stokes TM, Leonard CM. Central control of song in the canary, *Serinus canarius*. *J Comp Neurol*. 1976; 165:457–486. [PubMed: 1262540]
16. Coleman MJ, Vu ET. Recovery of impaired songs following unilateral but not bilateral lesions of nucleus uvaeformis of adult zebra finches. *J Neurobiol*. 2005; 63:70–89. [PubMed: 15685609]
17. Williams H, Vicario DS. Temporal patterning of song production: participation of nucleus uvaeformis of the thalamus. *J Neurobiol*. 1993; 24:903–912. [PubMed: 8228968]
18. Hahnloser RH, Kozhevnikov AA, Fee MS. An ultra-sparse code underlies the generation of neural sequences in a songbird. *Nature*. 2002; 419:65–70. [PubMed: 12214232]
19. Nottebohm F, Kelley DB, Paton JA. Connections of vocal control nuclei in the canary telencephalon. *J Comp Neurol*. 1982; 207:344–357. [PubMed: 7119147]
20. Thompson SM, Masukawa LM, Prince DA. Temperature dependence of intrinsic membrane properties and synaptic potentials in hippocampal CA1 neurons in vitro. *J Neurosci*. 1985; 5:817–824. [PubMed: 3973697]
21. Sabatini BL, Regehr WG. Timing of synaptic transmission. *Annu Rev Physiol*. 1999; 61:521–542. [PubMed: 10099700]
22. Volgushev M, Vidyasagar TR, Chistiakova M, Eysel UT. Synaptic transmission in the neocortex during reversible cooling. *Neuroscience*. 2000; 98:9–22. [PubMed: 10858607]
23. Volgushev M, Vidyasagar TR, Chistiakova M, Yousef T, Eysel UT. Membrane properties and spike generation in rat visual cortical cells during reversible cooling. *J Physiol*. 2000; 522(Pt 1): 59–76. [PubMed: 10618152]
24. Swadlow HA, Waxman SG, Weyand TG. Effects of variations in temperature on impulse conduction along nonmyelinated axons in the mammalian brain. *Exp Neurol*. 1981; 71:383–389. [PubMed: 7449905]
25. Lomber SG. The advantages and limitations of permanent or reversible deactivation techniques in the assessment of neural function. *J Neurosci Methods*. 1999; 86:109–117. [PubMed: 10065980]
26. Kalil RE, Chase R. Corticofugal influence on activity of lateral geniculate neurons in the cat. *J Neurophysiol*. 1970; 33:459–474. [PubMed: 4314735]
27. Katz PS, Sakurai A, Clemens S, Davis D. Cycle period of a network oscillator is independent of membrane potential and spiking activity in individual central pattern generator neurons. *J Neurophysiol*. 2004; 92:1904–1917. [PubMed: 15115787]
- **28. Tang LS, Goeritz ML, Caplan JS, Taylor AL, Fisek M, Marder E. Precise temperature compensation of phase in a rhythmic motor pattern. *PLoS Biol*. 2010; 8 The authors examine how the dynamics in interacting parts of the stomatogastric ganglion of the crab can be ‘temperature compensated’ to achieve robust operation over a wide range of temperatures and network oscillation frequencies.
29. Arbas EA, Calabrese RL. Rate modification in the heartbeat central pattern generator of the medicinal leech. *J Comp Physiol A*. 1984; 155:783–794.
- *30. Pires A, Hoy RR. Temperature coupling in cricket acoustic communication. II. Localization of temperature effects on song production and recognition networks in *Gryllus firmus*. *J Comp Physiol A*. 1992; 171:79–92. One of the first studies to use focal temperature changes to localize brain dynamics underlying a behavior. When heating was applied only to the thorax of a singing cricket, the speed of the song increased. When heating was applied only to the head, no change in song speed was detected. [PubMed: 1403993]
31. Bauer M, von Helversen O. Separate localization of sound recognizing and sound producing neural mechanisms in a grasshopper. *J Comp Physiol A*. 1987; 161:95–101.
32. Yamaguchi A, Gooler D, Herrold A, Patel S, Pong WW. Temperature-dependent regulation of vocal pattern generator. *J Neurophysiol*. 2008; 100:3134–3143. [PubMed: 18829853]
33. Long MA, Fee MS. Using temperature to analyse temporal dynamics in the songbird motor pathway. *Nature*. 2008; 456:189–194. [PubMed: 19005546]
34. Aronov D, Fee MS. Analyzing the dynamics of brain circuits with temperature: Design and implementation of a miniature thermoelectric device. *J Neurosci Methods*. 2011; 197:32–47. [PubMed: 21291909]
35. Amari S. Learning patterns and pattern sequences by self-organizing nets of threshold elements. *IEEE Transactions on Computers*. 1972; c-21:1197–1206.


36. Abeles, M. *Corticongs: Neural Circuits of the Cerebral Cortex*. Cambridge University Press; 1991.
37. Li M, Greenside H. Stable propagation of a burst through a one-dimensional homogeneous excitatory chain model of songbird nucleus HVC. *Phys Rev E Stat Nonlin Soft Matter Phys*. 2006; 74:011918. [PubMed: 16907138]
38. Jin DZ, Ramazanoglu FM, Seung HS. Intrinsic bursting enhances the robustness of a neural network model of sequence generation by avian brain area HVC. *J Comput Neurosci*. 2007; 23:283–299. [PubMed: 17440800]
39. Long MA, Jin DZ, Fee MS. Support for a synaptic chain model of neuronal sequence generation. *Nature*. 2010; 468:394–9. [PubMed: 20972420]
40. Foster DJ, Wilson MA. Hippocampal theta sequences. *Hippocampus*. 2007; 17:1093–1099. [PubMed: 17663452]
41. Buzsaki G. Two-stage model of memory trace formation: a role for “noisy” brain states. *Neuroscience*. 1989; 31:551–570. [PubMed: 2687720]
42. Mehta MR, Lee AK, Wilson MA. Role of experience and oscillations in transforming a rate code into a temporal code. *Nature*. 2002; 417:741–746. [PubMed: 12066185]
43. Henneman E. Relation between size of neurons and their susceptibility to discharge. *Science*. 1957; 126:1345–1347. [PubMed: 13495469]
44. Giocomo LM, Hasselmo ME. Time constants of h current in layer ii stellate cells differ along the dorsal to ventral axis of medial entorhinal cortex. *J Neurosci*. 2008; 28:9414–9425. [PubMed: 18799674]
- *45. Padmanabhan K, Urban NN. Intrinsic biophysical diversity decorrelates neuronal firing while increasing information content. *Nat Neurosci*. 2010; 13:1276–1282. The authors investigate the role of intrinsic excitability in neural circuit function. They measure biophysical parameters in single miral cells *in vitro* and find that biophysical heterogeneity helped to actively decorrelate neurons and increase information processing capacity. [PubMed: 20802489]
46. Solis MM, Perkel DJ. Rhythmic activity in a forebrain vocal control nucleus *in vitro*. *J Neurosci*. 2005; 25:2811–2822. [PubMed: 15772341]
- *47. Lee AK, Manns ID, Sakmann B, Brecht M. Whole-cell recordings in freely moving rats. *Neuron*. 2006; 51:399–407. The first demonstration intracellular recordings in a freely moving animal. The authors describe a head-mounted microdrive that was used to carry out intracellular patch recordings in cortical neurons of the rat. [PubMed: 16908406]
48. Epsztein J, Brecht M, Lee AK. Intracellular Determinants of Hippocampal CA1 Place and Silent Cell Activity in a Novel Environment. *Neuron*. 2011; 70:109–120. [PubMed: 21482360]
49. Buonomano DV, Maass W. State-dependent computations: spatiotemporal processing in cortical networks. *Nat Rev Neurosci*. 2009; 10:113–125. [PubMed: 19145235]
- *50. Branco T, Clark BA, Hausser M. Dendritic discrimination of temporal input sequences in cortical neurons. *Science*. 2010; 329:1671–1675. This paper shows how dynamics arising from dendritic impedance gradients and nonlinear activation of NMDA-type glutamate receptors leads to temporal sensitivity of a neuron to specific temporal patterns of synaptic input on the dendrites. [PubMed: 20705816]
- *51. Ganguli S, Bisley JW, Roitman JD, Shadlen MN, Goldberg ME, Miller KD. One-dimensional dynamics of attention and decision making in LIP. *Neuron*. 2008; 58:15–25. This paper suggests that the timecourse of changes in the locus of attention after the presentation of distractors in a visual task is the result of low-dimensional dynamics in LIP circuitry. It is suggested that recurrent excitation in randomly connected cortical networks leads to a single slow timescale that could control brain dynamics in a range of behaviors. [PubMed: 18400159]
52. Churchland MM, Cunningham JP, Kaufman MT, Ryu SI, Shenoy KV. Cortical preparatory activity: representation of movement or first cog in a dynamical machine? *Neuron*. 2010; 68:387–400. [PubMed: 21040842]
- **53. Harvey CD, Collman F, Dombeck DA, Tank DW. Intracellular dynamics of hippocampal place cells during virtual navigation. *Nature*. 2009; 461:941–946. This study describes whole-cell patch recordings of hippocampal neurons in a behaving mouse as it walks on a floating ball and navigates through a virtual environment. The direct measurement of intracellular membrane

- potentials allowed a test of different models of the dynamics that underlie the phenomenon of theta phase precession in place cells. [PubMed: 19829374]
54. O'Keefe J, Recce ML. Phase relationship between hippocampal place units and the EEG theta rhythm. *Hippocampus*. 1993; 3:317–330. [PubMed: 8353611]
 - *55. Fries P. Neuronal gamma-band synchronization as a fundamental process in cortical computation. *Annu Rev Neurosci*. 2009; 32:209–224. It is argued that gamma-band synchronization plays a fundamental role in regulating the communication between multiple interconnected cortical areas. The author also envisions a fundamental role for oscillatory brain dynamics in driving and controlling cortical computations. [PubMed: 19400723]
 - *56. Raichle ME, MacLeod AM, Snyder AZ, Powers WJ, Gusnard DA, Shulman GL. A default mode of brain function. *Proc Natl Acad Sci U S A*. 2001; 98:676–682. This work uses fMRI to observe spontaneous patterns of activity in the awake 'resting' human brain. The data suggest the existence of an organized, baseline 'default mode' of brain function. These spontaneous patterns of brain activity, more recently associated with 'stimulus-independent thought,' presumably involve complex internal dynamics in brain circuits. [PubMed: 11209064]
 - *57. Rivlin-Etzion M, Marmor O, Heimer G, Raz A, Nini A, Bergman H. Basal ganglia oscillations and pathophysiology of movement disorders. *Curr Opin Neurobiol*. 2006; 16:629–637. Parkinsonian tremor, one of the most disruptive symptoms of Parkinson's Disease, is associated with pathological synchronous oscillations in basal ganglia circuits and is an example of brain dynamics gone awry. This review points out that the mechanisms underlying the tremor are unknown, and that the tremor may be a compensatory response of BG circuits for other deficits. [PubMed: 17084615]
 58. Boyden ES, Zhang F, Bamberg E, Nagel G, Deisseroth K. Millisecond-timescale, genetically targeted optical control of neural activity. *Nat Neurosci*. 2005; 8:1263–1268. [PubMed: 16116447]
 59. Zhang F, Wang LP, Brauner M, Liewald JF, Kay K, Watzke N, Wood PG, Bamberg E, Nagel G, Gottschalk A, et al. Multimodal fast optical interrogation of neural circuitry. *Nature*. 2007; 446:633–639. [PubMed: 17410168]
 60. Huber D, Petreanu L, Ghitani N, Ranade S, Hromadka T, Mainen Z, Svoboda K. Sparse optical microstimulation in barrel cortex drives learned behaviour in freely moving mice. *Nature*. 2008; 451:61–64. [PubMed: 18094685]
 61. Lin D, Boyle MP, Dollar P, Lee H, Lein ES, Perona P, Anderson DJ. Functional identification of an aggression locus in the mouse hypothalamus. *Nature*. 2011; 470:221–226. [PubMed: 21307935]
 62. Hwang RY, Zhong L, Xu Y, Johnson T, Zhang F, Deisseroth K, Tracey WD. Nociceptive neurons protect *Drosophila* larvae from parasitoid wasps. *Curr Biol*. 2007; 17:2105–2116. [PubMed: 18060782]
 63. Ciochi S, Herry C, Grenier F, Wolff SB, Letzkus JJ, Vlachos I, Ehrlich I, Sprengel R, Deisseroth K, Stadler MB, et al. Encoding of conditioned fear in central amygdala inhibitory circuits. *Nature*. 2010; 468:277–282. [PubMed: 21068837]
 64. Stirman JN, Crane MM, Husson SJ, Wabnig S, Schultheis C, Gottschalk A, Lu H. Real-time multimodal optical control of neurons and muscles in freely behaving *Caenorhabditis elegans*. *Nat Methods*. 2011; 8:153–158. [PubMed: 21240278]
 65. Kravitz AV, Freeze BS, Parker PR, Kay K, Thwin MT, Deisseroth K, Kreitzer AC. Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. *Nature*. 2010; 466:622–626. [PubMed: 20613723]
 66. Schoonheim PJ, Arrenberg AB, Del Bene F, Baier H. Optogenetic localization and genetic perturbation of saccade-generating neurons in zebrafish. *J Neurosci*. 2010; 30:7111–7120. [PubMed: 20484654]
 67. Lerchner W, Xiao C, Nashmi R, Slimko EM, van Trigt L, Lester HA, Anderson DJ. Reversible silencing of neuronal excitability in behaving mice by a genetically targeted, ivermectin-gated Cl⁻ channel. *Neuron*. 2007; 54:35–49. [PubMed: 17408576]
 68. Tan EM, Yamaguchi Y, Horwitz GD, Gosgnach S, Lein ES, Goulding M, Albright TD, Callaway EM. Selective and quickly reversible inactivation of mammalian neurons in vivo using the *Drosophila* allatostatin receptor. *Neuron*. 2006; 51:157–170. [PubMed: 16846851]

69. Hagglund M, Borgius L, Dougherty KJ, Kiehn O. Activation of groups of excitatory neurons in the mammalian spinal cord or hindbrain evokes locomotion. *Nat Neurosci.* 2010; 13:246–252. [PubMed: 20081850]
70. Zhang Y, Narayan S, Geiman E, Lanuza GM, Velasquez T, Shanks B, Akay T, Dyck J, Pearson K, Gosgnach S, et al. V3 spinal neurons establish a robust and balanced locomotor rhythm during walking. *Neuron.* 2008; 60:84–96. [PubMed: 18940590]
- **71. Gosgnach S, Lanuza GM, Butt SJ, Saueressig H, Zhang Y, Velasquez T, Riethmacher D, Callaway EM, Kiehn O, Goulding M. V1 spinal neurons regulate the speed of vertebrate locomotor outputs. *Nature.* 2006; 440:215–219. The authors investigated the role of the V1 population in a locomotor central pattern generator of the spinal cord. Through a variety of methods, they show that V1 neurons regulate the dynamics of locomotor rhythms. [PubMed: 16525473]

Highlights

> Some circuits in the brain drive the timing and ordering of behavior, others follow. > Existing techniques are not suited to localizing and characterizing circuit dynamics. > Localized mild cooling of brain areas can be used to detect dynamics. > Models of circuit dynamics can also be tested with intracellular recording and current injection.

**FIGURE 1.**

Sequential activity in the songbird motor pathway. A, Block diagram of brain regions involved in the production of song. Abbreviations: Uva, nucleus Uvaeformis; HVC, used as a proper name; RA, robust nucleus of the archistriatum; MN, brainstem motor nucleus. B, Raster plot of the spiking patterns of eight RA-projecting HVC neurons recorded during singing. Each row of tick marks shows spikes generated during one rendition of the song; roughly ten renditions are shown for each neuron. The neurons were identified by antidromic stimulation from RA. The song spectrogram is shown at top. Note that each RA-projecting HVC neuron bursts reliably at a single precise time in the song.





FIGURE 2.

Localizing dynamics underlying sequence generation in HVC. A, Sequential bursts within HVC drive precise sequences of activity in the downstream motor pathway that produce song behavior. B, In Model 1 sequences are generated by circuitry within HVC. Slowing circuit dynamics in HVC should therefore slow down the sequence of bursts, thus slowing the song. Grey and blue arrows indicate speed of control and cooled burst sequence, respectively. Bars below indicate duration of control and manipulated song syllables. C, In Model 2 the dynamics underlying song timing reside upstream of HVC, for example in Uva. HVC bursts are triggered by timing signals from Uva. Slowing circuit dynamics in HVC might increase the latency of the HVC response to Uva input but this latency increase will be the same for every burst. (Increased latency represented by rightward shift of blue arrow.) Thus, the number of bursts per unit time will not change and the song will not slow down.

**FIGURE 3.**

Slowing dynamics with local temperature change. A, Schematic of the zebra finch brain highlighting the motor pathway and showing the placement of the thermoelectric cooling device over HVC. B, Bidirectional temperature changes at a variety of depths under the surface of the thermoelectric probe, as a function electrical current through the device. Note that the temperature changes are localized to within 1-2mm of the cooling device: The distance to the center of HVC is ~ 0.5 mm and to the center of RA is ~ 4 mm. C, Cooling HVC produces a slowing of the song across all timescales. Shown are song spectrograms from a bird with HVC at normal brain temperatures (top) and during cooling of HVC by 8° C (middle). At bottom is a linearly warped image of the normal song spectrogram.




FIGURE 4.

Testing circuit dynamics during natural behaviors with intracellular current injection and recording. **A**, In the chain model of sequence generation, neurons activate each other sequentially through sparse feed-forward synaptic connections. This model predicts that intracellular current injection should have little effect on burst timing. Strong hyperpolarizing currents (red) might suppress spiking, but there should be a subthreshold depolarizing potential at the time of the burst. **B**, In the ramp-to-threshold model, neurons are activated by a global ramp of excitation. The time at which each neuron spikes is set by its intrinsic excitability. This model predicts that burst timing should be delayed by hyperpolarizing current (red) and advanced by depolarizing current injection (green). **C**,

Motorized microdrive for intracellular recording and current injection in unrestrained, freely behaving animals. D, Intracellular recording of a single RA-projecting HVC neuron during singing. The membrane potential is aligned to the song motif. Traces are shown at low and high voltage gains (top and bottom, respectively). Note the absence of slow membrane potential ramping prior to burst onset. E, Consistent with the chain model, burst timing was unaffected by steady current injection. Note the depolarizing potential at the time of the burst in the most hyperpolarized condition (bottom).